## **Allergy and Anaphylaxis Emergency Plan**

		<del></del>		
Name:	Date of Birth:		Weight:	lbs / kg
Date of Plan:	Age:			
ALLERGIES:				
Child has asthma: yes / no (if yes, higher chance of a severe reaction) Child has had anaphylaxis: yes / no (if yes, higher chance of a severe reaction) Child may carry medicine: yes / no Child may give him/herself medicine: yes / no (if child refuses, an adult must give medicine)				Attach child's
☐ The "Always-Epinephrine" Option: If checked, give epinephrine immediately, if the child has ANY symptom (mild or severe) after a sting or eating a food listed above.  (Option advised for those schools where a nurse is not always present.)				
**IF IN DOUBT, GIVE EPINEPHRINE! ANAPHYLAXIS is a potentially life-threatening, severe allergic reaction				
For SEVERE Allergy or Anaphylax What to look for:  If child has ANY of these symptoms food or having a sting, give epineph  ➤ Breathing: trouble breathing, whe Throat: tight or hoarse throat, trous or speaking  ➤ Brain: confusion, agitation, dizzing unresponsiveness  ➤ Gut: severe stomach pain, vomitite Mouth: swelling of lips or tongue breathing  ➤ Skin: face color is pale or blue, many redness over body	after eating a arrine eeze, cough uble swallowing ness, fainting, ing, diarrhea that affects	<ul> <li>Tell rescue squa</li> <li>Stay with child and</li> <li>Call parents</li> <li>Give a second of worsen or do not</li> <li>Keep child lying of trouble breathing</li> <li>Give other medicin</li> </ul>	e right away! Note to be with epinephrine d when epinephrine : dose of epinephrine is get better in 5 minus on back. If the child g, keep child lying or	e was given  e if symptoms utes vomits or has n their side e, inhaler) if
For MILD Allergic Reaction What to look for: If child has mild symptoms, or no syr sting or ingestion of the food is suspending and monitor the child. Mild symptoms may include:  > Skin: a few hives, mild rash, mild.  Mouth/nose/eyes: itching, rubbing.  Gut: mild stomach pain, nausea of the child has more than one area affected, give epinephrine	ected, give swelling, OR g, sneezing, OR or discomfort	Give Antihistamine What to do:  1. Give antihistamine 2. If in doubt, give of 3. Call parents 4. Watch child close 5. If symptoms work SEVERE Allergy of	e if prescribed epinephrine ly for 4 hours sen, give epinephi	
Medicine/Doses Epinephrine (intramuscular in thigh): Antihistamine (by mouth): □ Diphenh Other medications: □ Albuterol 2-4 p	nydramine uffs □ other:	_mg(ml)□ Othe		
PROVIDER Signature	Date N	lame (printed)	Phone	FAX

Reviewed by school nurse: \_\_\_\_\_\_ Date: \_\_\_\_\_

I authorize the school to follow Plan and contact the Health Care Provider, and release the school district and personnel from civil liability

Name (printed) Phone

PARENT/GUARDIAN Signature Date

## **Allergy and Anaphylaxis Emergency Plan**

Child's name:	Date of Plan:		
Additional Instructions:			
Contacts			
Doctor name (print):	Office Phone: ( ) -		
Office Address:	Office Four (		
	Phone:		
Parent/Guardian name (print) :	Phone:		
Other Emergency Contacts			
Name/Relationship:	Phone:		
Name/Relationship:			
Decidence de la constant de la const	5.1		
Reviewed by school nurse:	Date:		